




ARKLE Veterinary Care, LLC
1020 Concord Rd., SE • Smyrna, GA 30080
Phone: (770) 435-6700 • Fax: (770) 434-4863
www.arklevetcare.com • arklevetcare@bellsouth.net

KENNEL COUGH

What is Kennel Cough?

Kennel Cough is a broad term covering any infectious or contagious condition of dogs where coughing is one of the major clinical signs. The term *tracheobronchitis* describes the location of the infection in the “windpipe” or trachea and bronchial tubes. Several viruses and bacteria can cause kennel cough, often at the same time. These include adenovirus type-2 (distinct from the adenovirus type 1 that causes *infectious hepatitis*), parainfluenza virus, and the bacterium *Bordetella bronchiseptica*. Because the infection spreads when dogs are housed together, it is often seen soon after dogs have been in kennels, hence the name “kennel cough”.



What are the clinical signs of kennel cough other than coughing?

Clinical signs may be variable. It is often a mild disease, but the cough may be chronic, lasting for several weeks in some cases. Common clinical signs include a loud cough often describe as a “goose honk”, runny eyes and nose, swollen tonsils, wheezing, lack of appetite and depressed behavior. Most cases of infectious tracheobronchitis have a demonstrable or elicitable cough that occurs when the throat is rubbed or palpated.

What is the treatment for infectious tracheobronchitis?

There is no specific treatment for the viral infections, but many of the more severe signs are due to bacterial involvement, particularly *Bordetella bronchiseptica*. Antibiotics are useful against this bacterium, although some antibiotic resistance has been reported. Some cases require prolonged treatment, but most infections resolve within one to three weeks. Mild clinical signs may linger even when the bacteria have been eliminated.

How can I prevent my dog contracting Kennel Cough?

Most vaccination programs your veterinarian will recommend contain adenovirus and parainfluenza. Bordetella vaccination is also highly recommended for dogs that are boarded, groomed or interact with other dogs in areas such as dog parks.

How effective are these vaccines?

Immunity, even if the dog has experienced a natural infection, is neither solid nor long-lasting. We cannot expect vaccines to do much better. Since immunity varies with the circumstances, we recommend vaccinating your pet every 6 months with the intranasal vaccine. There is an injectable vaccine, but immunity will develop slower from this than the intranasal form. Some kennel facilities require a booster vaccination shortly before boarding. We recommend you have your pet boosted 2 weeks before boarding unless he/she is getting regular 6 month boosters.

How are the Bordetella vaccines administered?

Bordetella vaccination is given either by injection or intra-nasal route. Intra-nasal refers to the liquid vaccine administered as nose drops. This allows local immunity to develop on the mucous membranes of the nose, throat and windpipe where the infectious agents first attack.